





General production advice warepotatoes

versie 2009 - 02

- * Very early
- * Bold tubers
- * Uniform size
- * Robust variety
- * Early harvest of bold tubers





Characteristics

Cooking type AB

Maturity Middle early
Yield mature Rather high
Tuber size Very large
Tuber shape Round-oval / Oval

Flesh colour Yellow
Skin colour Yellow
Berries Few berries

Dormancy period Average
Emergence Normal
Sencor sensitivity Rather sensitive

Sencor sensitivity Rather se Foliage development Good

Internal bruising
Little Potato disorder
Dry matter content

Good
Not sensitive
Rather sensitive
18,4

UWW 333

Wart disease

PCN Resistance Resistant: Ro1,4

Slightly susceptible: Ro2/3

Susceptible: Ro5 Resistant: fysio 1

Foliage Blight Slightly susceptible Tuber Blight Susceptible

Common scab
Powdery scab
PVYn
Resistant
Very susceptible
Susceptible

Yntn tuber tolerance Slightly susceptible

Plant populations

Advice planting densities (75 cm. ridge distance).

28/35mm-approx. 49.500 pl./ha. Plant. dens. \pm 27 cm. 35/50mm-approx. 44.500 pl./ha. Plant. dens. \pm 30 cm. 50/55mm-approx. 40.500 pl./ha. Plant. dens. \pm 33 cm.

Fertilizer

Adapt fertilization to soil analysis.

Broad Nitrogen dose (split) is desirable. Advice approx. 20 kg N/ha inclusive soil supply. The best results are being obtained with an initial dose of 200 kg.

Potassium- and Phosphate fertilization as per normal advice.

To be able to harvest early, organic fertilization is NOT advised.





General production advice warepotatoes

Pre-treatment and planting



- Store in cool conditions till springtime and then warm up thoroughly.
- A heat treatment is desirable.
- Pre sprouting is not necessary, variety can be planted good with small white sprouts.
- Prevent broken sprouts during planting.
- Do not plant in too cold soil, Carlita is susceptible to Little Potato Disorder.
- Plant somewhat above the soil surface.
- The best results are being obtained on lighter soils, not too susceptible to common Scab.

Growing attentionpoints



- The initial development is slow, followed by a smooth leaf development and a medium high plant with good cover.
- Tuberization is early, tuber numbers rather low. Tubers will be bold and a uniform size.
- Irrigation can increase the tuber numbers .
- Take care of uniform growth to prevent growth cracks.
- For an early variety Carlita is not very susceptible to Leaf Blight.
- Good drought resistance.

Haulmkilling and harvest



- Carlita allows an early harvest.
- The variety is somewhat susceptible to mechanical damage.
- Under normal conditions, harvest should not cause any problems.

Storage



- Good dormancy and good storage.
- For long storage mechanical cooling of 4 °C will give good results.
- Less susceptible to Fusarium.