




General production advice warepotatoes

version 2009 - 01

- * For an early variety good yield
- * Bold tubers
- * Strong
- * Excellent taste
- * Regular shape





Characteristics

Cooking type Maturity Yield mature Tuber size Tuber shape Number of tubers Flesh colour Skin colour Berries

Dormancy period Emergence Sencor sensitivity Foliage development Internal bruising Little Potato disorder Dry matter content UWW

PCN Resistance

Wart disease

Spraing Foliage Blight Tuber Blight Common scab Powdery scab PVYn Yntn tuber tolerance Early Good Very large Round-oval / Oval Low tuber number Light yellow Yellow No berries

Average Normal Sensitive Good Not sensitive Rather sensitive 19,3 351

Susceptible: Ro1,2/3,4,5 Pa2,3 Resistant: fysio 1

Resistant Susceptible Slightly susceptible Susceptible Susceptible Slightly susceptible

Plant populations

Advice planting distance (row distance 75 cm.)

28/35 mm - ca. 51.000 pl./ha. Pl. distance \pm 26 cm 35/50 mm - ca. 47.500 pl./ha. Pl. distance \pm 28 cm 50/55 mm - ca. 45.500 pl./ha. Pl. distance \pm 30 cm

Fertilizer

An ample Nitrogen fertilisation is advised. Advise about 275 kg N/ha. including soil supply. A good Potassium and Phosphorus fertilisation can increase the yield.

HZPC takes no responsibility for any harmful consequences that might eventually occur when using this information.





General production advice warepotatoes

Pre-treatment and planting	
	 For normal sprouting, a heat treatment is usually not necessary. Sensitive to little potato disorder. Do not desprout more than once and don't plant too early in too cold soil.
Growing attentionpoints	
	 Sensitive to Sencor. Emergence is rather quick. Good foliage development and coverage. Tuberisation is early. Tubers will quickly mature to a bold size. Begular and timely spraying against late Blight (Phytophthora) is important.

- Regular and timely spraying against late Blight (Phytophthora) is important.

Haulmkilling and harvest



- Little sensitivity to mechanical damage. - Not sensitive to bruising.

Storage



- The dormancy is moderately long, the storability rather good.
 Can be stored in a mechanical cold store at 4℃.

